DESY-98-106
hep-ph/9808329

arXiv:hep-ph/9808329v1 14 Aug 1998

Charmless Nonleptonic Two-Body B Decays in the Factorization Approach

Cai-Dian Lü[1]
II Institut für Theoretische Physik, Universität Hamburg, D - 22761 Hamburg, Germany

Abstract

We calculate the two-body nonleptonic B decays using the factorization method. The recent measured decays by CLEO Collaboration can be explained in the factorization approach. We propose a number of ratios of branching ratios to determine the effective coefficients and the form factors.

Invited talk given at the
6th International Euroconference QCD 98
Montpellier, France, 2–8 July 1998
To appear in the Proceedings

[1]Alexander von Humboldt research fellow.

Charmless nonleptonic two-body B decays in the factorization approach

C.-D. Lü
II Institut für Theoretische Physik, Universität Hamburg, 22761 Hamburg, Germany†

We calculate the two-body nonleptonic B decays using the factorization method. The recent measured decays by CLEO Collaboration can be explained in the factorization approach. We propose a number of ratios of branching ratios to determine the effective coefficients and the form factors from experiments.

1. INTRODUCTION

In this report I represent works done in collaboration with A. Ali and G. Kramer [1].

Recently the CLEO Collaboration has measured a number of charmless nonleptonic B decays [2]. This has aroused great interest of studying these decays [3–5]. The charmless nonleptonic B decays provide the opportunity to study CP violation and a way of measuring of CKM matrix elements. Since they are rare decays, they are also useful for tests of the Standard Model and giving signals of possible new physics.

The calculation of nonleptonic decays involves the short-distance Lagrangian and the calculation of hadronic matrix elements which are model dependent. The short-distance QCD corrected Lagrangian is calculated to next-to-leading order. The popular method to calculate the hadronic matrix elements is using the factorization method where the matrix element is expressed as a product of two factors $\langle h_1 h_2|\mathcal{H}_{eff}|B\rangle = \langle h_1|J_1|B\rangle\langle h_2|J_2|0\rangle$.

This model works well for the tree level hadronic B and D decays using phenomenological values of a_1 and a_2 from experiments [6,7]. Does it also work for decays involving light mesons? In charmless decays, penguins play an important role. What is the effective number of color N_c in this category of decays? Here we use the factorization method to analyze the charmless nonleptonic decays of 76 channels and their charge conjugated decays. A number of ratios is proposed to test the factorization approach and measure the parameters a_i and CKM matrix elements.

†Alexander von Humboldt research fellow.

2. THEORETICAL FRAMEWORK

The effective Hamiltonian for the charmless nonleptonic B decays is

$$\mathcal{H}_{eff} = \frac{G_F}{\sqrt{2}}\left[V_{q'b}V_{q'q}^*\left(\sum_{i=1}^{10} C_i\,O_i + C_g O_g\right)\right], \quad (1)$$

where $q = d, s$ and $V_{q'q}$ denotes the CKM factors. The operators O_1, O_2 are current operators. The operators $O_3, \ldots, O_6$ are QCD penguin operators. $O_7, \ldots, O_{10}$ arise from electroweak penguin diagrams, which are suppressed by α/α_s. Only O_9 has a sizable value whose major contribution arises from the Z penguin. In practice, we work at $\mu = 2.5\ GeV$, in the naive dimensional regularization scheme.

For example, let us consider B meson decays to two pseudoscalar mesons. In the factorization approach, using the effective Hamiltonian, we write the required matrix element in its factorized form

$$\langle P_1 P_2|\mathcal{H}_{eff}|B\rangle = i\frac{G_F}{\sqrt{2}}V_{qb}V_{qq'}^* a_i f_{P_2}(m_B^2 - m_1^2)F_0^{B\to P_1}(m_2^2). \quad (2)$$

The dynamical details are coded in the quantities a_i, which we define as $a_i \equiv C_i^{eff} + C_j^{eff}/N_c$, where $\{i, j\}$ is any of the pairs $\{1,2\}$, $\{3,4\}$, $\{5,6\}$, $\{7,8\}$ or $\{9,10\}$. In practice, N_c is treated as a phenomenological parameter to include the non-factorized color-octet contributions.

The QCD and electroweak penguins are also present in the charmless decays of B mesons. However, if the amplitude is still dominated by the tree amplitude, the BSW-classification can be applied as before [6]. Class-I decays are color favored, whose matrix elements are proportional to

Table 1
$B \to h_1 h_2$ Branching Ratios (in units of 10^{-6}).

Channel	$N_c = 2$	3	∞	Exp.
$B^0 \to \pi^+\pi^-$	9.0	10.0	12	< 8.4
$B^+ \to \pi^+\pi^0$	6.8	5.4	3.0	< 16
$B^+ \to K^+\pi^0$	9.4	10	12	$15 \pm 4 \pm 3$
$B^0 \to K^+\pi^-$	14	15	18	$14 \pm 3 \pm 2$
$B^+ \to K^+\eta'$	21	25	35	$74^{+8}_{-13} \pm 9$
$B^0 \to K^0\eta'$	20	25	35	$59^{+18}_{-16} \pm 9$
$B^+ \to \pi^+K^0$	14	16	22	$14 \pm 5 \pm 2$
$B^0 \to \rho^+\pi^-$	21	23	28	< 88
$B^+ \to \rho^+\pi^0$	14	13	11	< 77
$B^+ \to K^{*0}\pi^+$	5.6	6.9	10	< 39
$B^+ \to \omega K^+$	3.2	0.25	11	$15^{+7}_{-6} \pm 2$
$B^0 \to \rho^+\rho^-$	18	20	24	< 2200

a_1. Class-II decays are color suppressed, whose matrix elements are proportional to a_2. Class-III decays are proportional to $a_1 + r a_2$.

For the penguin-dominated decays, we introduce two new classes: Class-IV decays involve one or more of the dominant penguin coefficients a_4, a_6 and a_9. Class-V decays are decays with strong N_c-dependent coefficients a_3, a_5, a_7 and a_{10}.

Class-I and Class-IV decays have relatively large branching ratios of the order of 10^{-5} and stable against variation of N_c. Class-III decays are mostly stable, except some $B \to PV$ decays. Class-II and Class-V decays are rather unstable against variation of N_c. Many of them may receive significant contribution from the annihilation diagrams and/or soft final state interactions.

3. NUMERICAL CALCULATIONS

The CKM matrix is expressed in terms of the Wolfenstein parameters. We take $A = 0.81$, $\lambda = \sin\theta_C = 0.2205$, $\rho = 0.12$, $\eta = 0.34$. This choice of ρ, η corresponds to CKM triangles: $\alpha = 88.3^\circ$, $\beta = 21.1^\circ$, $\gamma = 70.6^\circ$. At scale $\mu = 2.5$GeV, we use current masses $m_b = 4.88$ GeV, $m_c = 1.5$ GeV, $m_s = 122$ MeV, $m_d = 7.6$ MeV and $m_u = 4.2$ MeV. The form factors are taken from ref. [6,1] indicated as BSW model and Lattice QCD/QCD sum rule. The decay constants are shown in ref. [1].

We select a few decay branching ratios shown in Table 3. For the whole list of the 76 channels, see ref. [1]. In this table, it is easy to see that $B^0 \to K^+\pi^-$, $B^+ \to K^+\eta'$, $B^0 \to K^0\eta'$, $B^+ \to \pi^+K^0$, measured by CLEO, are well explained by factorization. The experiment of $B^+ \to K^+\eta'$ favors a small value of $\xi = 1/N_c$.

The CLEO experiments measured a combined channel $\mathcal{B}(B^+ \to \pi^0 h^+) = (1.6^{+0.6}_{-0.5} \pm 0.4) \times 10^{-5}$, $(h^+ = \pi^+, K^+)$. The measurements agree with experiments for all possible values of ξ. The CLEO Collaboration also measured $\mathcal{B}(B^+ \to \omega h^+) = (2.5^{+0.8}_{-0.7} \pm 0.3) \times 10^{-5}$. This suggests values of $\xi \simeq 0$ and $\xi \geq 0.5$. However, since $B^+ \to \omega K^+$ is a Class V decay and $B^+ \to \omega\pi^+$ is a class III decays, they are unstable. Thus it is too early to draw definite conclusions. We expect that $B^0 \to \pi^+\pi^-$, $\rho^+\pi^-$, $\rho^+\rho^-$ and $B^+ \to \pi^+\pi^0$, $K^+\pi^0$ will be observed in the next round of experiments at CLEO and at B factories.

4. USEFUL RATIOS

We start with the ratios independent of the effective coefficients a_i. Neglecting the small QCD penguin contribution and the very small difference in phase space, we get the relations:

$$P_2 \equiv \frac{\mathcal{B}(B^0 \to \pi^-\pi^+)}{\mathcal{B}(B^0 \to \rho^+\pi^-)} \simeq \left(\frac{f_\pi F_0^{B\to\pi}}{f_\rho F_1^{B\to\pi}}\right)^2,$$

$$P_3 \equiv \frac{\mathcal{B}(B^0 \to \pi^+\rho^-)}{\mathcal{B}(B^0 \to \rho^+\pi^-)} \simeq \left(\frac{f_\pi A_0^{B\to\rho}}{f_\rho F_1^{B\to\pi}}\right)^2. \quad (3)$$

These two ratios P_2 and P_3 can measure $F_0^{B\to\pi}/F_1^{B\to\pi}$ and $A_0^{B\to\rho}/F_1^{B\to\pi}$, respectively.

$$P_4 \equiv \frac{\mathcal{B}(B^+ \to \pi^+\pi^0)}{\mathcal{B}(B^+ \to \rho^+\rho^0)} \simeq \frac{f_\pi^2}{f_\rho^2}\frac{|F_1^{B\to\pi}|^2}{(1+x)|A_1^{B\to\rho}|^2},$$

$$P_5 \equiv \frac{\mathcal{B}(B^0 \to \pi^-\pi^+)}{\mathcal{B}(B^0 \to \rho^-\rho^+)} \simeq \frac{f_\pi^2}{f_\rho^2}\frac{|F_1^{B\to\pi}|^2}{(1+x)|A_1^{B\to\rho}|^2}, \quad (4)$$

where $x = m_\rho/m_B$. The relation $P_4 = P_5$ provides a test of factorization.

$$P_1 \equiv \frac{\mathcal{B}(B^0 \to \rho^+\pi^-)}{\mathcal{B}(B^0 \to \rho^+\rho^-)} \simeq \frac{|F_1^{B\to\pi}|^2}{(1+x)|A_1^{B\to\rho}|^2},$$

$$P_6 \equiv \frac{\mathcal{B}(B^0 \to K^{*+}\pi^-)}{\mathcal{B}(B^0 \to K^{*+}\rho^-)} \simeq \frac{|F_1^{B\to\pi}|^2}{(1+x)|A_1^{B\to\rho}|^2},$$

Table 2
Values of P_i's using the BSW form factors and the lattice-QCD/QCD-sum rule form factors. The numbers in brackets are calculated using the approximate formulae derived in the text.

Ratio	BSW model	Lattice-QCD
P_1	1.19 [1.21]	1.27 [1.55]
P_2	0.43 [0.39]	0.43 [0.39]
P_3	0.28 [0.28]	0.27 [0.27]
P_4	0.49 [0.47]	0.53 [0.61]
P_5	0.52 [0.47]	0.55 [0.61]
P_6	1.11 [1.21]	1.19 [1.55]
P_7	1.11 [1.21]	1.19 [1.55]
P_8	1.08 [1.14]	0.99 [1.18]
P_9	1.09 [1.14]	0.99 [1.18]
P_{10}	1.01 [1.15]	0.92 [1.19]
P_{11}	1.01 [1.15]	0.92 [1.19]

$$P_7 \equiv \frac{\mathcal{B}(B^+ \to \pi^+ K^{*0})}{\mathcal{B}(B^+ \to \rho^+ K^{*0})} \simeq \frac{|F_1^{B\to\pi}|^2}{(1+x)|A_1^{B\to\rho}|^2}. \quad (5)$$

They are essentially determined by the ratios of the form factors $F_1^{B\to\pi}$ and $A_1^{B\to\rho}$. The relation $P_1 \simeq P_6 \simeq P_7$ can be a test of factorization.

$$P_8 \equiv \frac{\mathcal{B}(B^+ \to K^+ \bar{K}^{*0})}{\mathcal{B}(B^+ \to K^{*+} \bar{K}^{*0})},$$
$$P_9 \equiv \frac{\mathcal{B}(B^0 \to K^0 \bar{K}^{*0})}{\mathcal{B}(B^0 \to K^{*0} \bar{K}^{*0})},$$
$$P_{10} \equiv \frac{\mathcal{B}(B^+ \to K^+ \phi)}{\mathcal{B}(B^+ \to K^{*+} \phi)},$$
$$P_{11} \equiv \frac{\mathcal{B}(B^0 \to K^0 \phi)}{\mathcal{B}(B^0 \to K^{*0} \phi)}. \quad (6)$$

Ignoring the small phase space difference,

$$P_8 \simeq P_9 \simeq P_{10} \simeq P_{11} \simeq \frac{|F_1^{B\to K}|^2}{(1+y)|A_1^{B\to K^*}|^2},$$

where $y = m_{K^*}/m_B$. They are all proportional to the ratios of the form factors $F_1^{B\to K}$ and $A_1^{B\to K^*}$. The values of P_1-P_{11} calculated in our model are displayed in Table 2, together with values calculated using the approximate formulae. The two sets of numbers are quite close means that the approximations are working quite well.

In the following, we first give ratios to determine the effective coefficients a_1 and a_2

$$S_1 \equiv \frac{\mathcal{B}(B^0 \to \pi^+\pi^-)}{2\mathcal{B}(B^+ \to \pi^+\pi^0)} \simeq \left(\frac{a_1}{a_1+a_2}\right)^2,$$
$$S_2 \equiv \frac{\mathcal{B}(B^0 \to \rho^+\rho^-)}{2\mathcal{B}(B^+ \to \rho^+\rho^0)} \simeq \left(\frac{a_1}{a_1+a_2}\right)^2,$$
$$S_3 \equiv \frac{2\mathcal{B}(B^+ \to \rho^+\pi^0)}{\mathcal{B}(B^0 \to \rho^+\pi^-)} \simeq \left(1+\frac{1}{x}\frac{a_2}{a_1}\right)^2,$$
$$S_4 \equiv \frac{2\mathcal{B}(B^+ \to \pi^+\rho^0)}{\mathcal{B}(B^0 \to \pi^+\rho^-)} \simeq \left(1+x\frac{a_2}{a_1}\right)^2. \quad (7)$$

Similarly for penguin operators,

$$S_5 \equiv \frac{2\mathcal{B}(B^+ \to \pi^+\pi^0)}{\mathcal{B}(B^+ \to \pi^+ K^0)} \simeq \xi^2 \frac{f_\pi^2}{f_K^2}\left|\frac{a_1+a_2}{a_4+a_6R_5}\right|^2,$$
$$S_6 \equiv \frac{2\mathcal{B}(B^+ \to \rho^+\rho^0)}{\mathcal{B}(B^+ \to \rho^+ K^{*0})} \simeq \xi^2 \frac{f_\rho^2}{f_{K^*}^2}\left|\frac{a_1+a_2}{a_4}\right|^2,$$
$$S_7 \equiv \frac{\mathcal{B}(B^0 \to \pi^-\rho^+)}{\mathcal{B}(B^+ \to \pi^+ K^{*0})} \simeq \xi^2 \frac{f_\rho^2}{f_{K^*}^2}\left|\frac{a_1}{a_4}\right|^2,$$
$$S_8 \equiv \frac{\mathcal{B}(B^0 \to \rho^-\rho^+)}{\mathcal{B}(B^+ \to \rho^+ K^{*0})} \simeq \xi^2 \frac{f_\rho^2}{f_{K^*}^2}\left|\frac{a_1}{a_4}\right|^2,$$
$$S_9 \equiv \frac{\mathcal{B}(B^0 \to \pi^+\pi^-)}{\mathcal{B}(B^+ \to \pi^+ K^0)} \simeq \xi^2 \frac{f_\pi^2}{f_K^2}\left|\frac{a_1}{a_4+a_6R_5}\right|^2,$$

with $\xi = |V_{ub}V_{ud}^*|/|V_{tb}V_{ts}^*|$. We can use $S_3, ..., S_9$ to determine a_4 and a_6. The dominant contribution of the electroweak penguin amplitudes is proportional to a_9. It can be determined by the following two ratios.

$$S_{10} \equiv \frac{2\mathcal{B}(B^0 \to \rho^0 K^0)}{\mathcal{B}(B^+ \to \pi^+ K^{*0})} \simeq \frac{9}{4}\left|\frac{f_\rho F_1^{B\to K}}{f_{K^*}F_1^{B\to\pi}}\right|^2 \left|\frac{a_9}{a_4}\right|^2,$$
$$S_{11} \equiv \frac{2\mathcal{B}(B^0 \to \rho^0 K^0)}{\mathcal{B}(B^0 \to \rho^-\pi^+)} \simeq \frac{9}{4\xi^2}\left|\frac{f_\rho F_1^{B\to K}}{f_\pi A_0^{B\to\rho}}\right|^2 \left|\frac{a_9}{a_1}\right|^2.$$

It is difficult to quantitatively determine the other penguin coefficients which are smaller.

The ratio discussed by Fleischer and Mannel recently to constrain the CKM parameter γ [9],



Figure 1. S_{12} as a function of $\cos\gamma$. The dotted, dash-dotted and dashed curves correspond to $N_c = \infty$, $N_c = 3$ and $N_c = 2$, respectively. The horizontal lines are the CLEO ($\pm 1\sigma$) measurements of S_{12}.

is defined as

$$S_{12} \equiv \frac{\mathcal{B}(B^0 \to K^+\pi^-)}{\mathcal{B}(B^+ \to K^0\pi^+)} \simeq 1 - 2z_{12}\cos\delta_{12}\cos\gamma + z_{12}^2, \tag{8}$$

where δ_{12} is the strong phase, and z_{12} is defined in ref. [1]. The ratio S_{12} has the experimental value: $S_{12} = 1.0 \pm 0.4$. It is shown in Fig. 1. We see that the measurement of S_{12} can determine the CKM parameter $\cos\gamma$. Analogous to eqn. (8), we define

$$S_{13} \equiv \frac{\mathcal{B}(B^0 \to \pi^- K^{*+})}{\mathcal{B}(B^+ \to \pi^+ K^{*0})}$$
$$S_{15} \equiv \frac{\mathcal{B}(B^0 \to \rho^- K^{*+})}{\mathcal{B}(B^+ \to \rho^+ K^{*0})} \tag{9}$$

$S_{13} \simeq S_{15} \simeq 1 - 2z_{13}\cos\delta_{13}\cos\gamma + z_{13}^2$.

They can also be used to determine the $\cos\gamma$.

5. SUMMARY

The recently measured charmless nonleptonic B decay modes can be explained in the factorization framework. They show some preference for $\xi \leq 0.2$. A good fraction of the seventy six decay modes will be measured in the future providing a detailed test of the factorization approach.

We have put forward numerous proposals for ratios of branching ratios to determine the effective coefficients a_1, a_2, a_4, a_6 and a_9. We proposed also a number of ratios P_i, which will help in determining the form factors for the various decays considered here. The two-body nonleptonic decays provide potentially non-trivial constraints on the CKM parameters.

Finally, it is instructive to study direct and indirect CP violation in all the two-body nonleptonic B decays considered here [8].

REFERENCES

1. A. Ali, G. Kramer, C.-D. Lü, DESY-98-041, hep-ph/9804363, to appear in Phys. Rev. D.
2. M. Artuso et al. (CLEO Collaboration), preprint CONF 98-20; B.H. Behrens et al. (CLEO Collaboration), preprint CLEO CONF 98-09.
3. A. Ali, C. Greub, Phys. Rev **D57**, 2996 (1998); A. Ali, J. Chay, C. Greub and P. Ko, Phys. Lett. B424, 161 (1998).
4. N.G. Deshpande, B. Dutta, Sechul Oh, Phys. Rev. D57, 5723 (1998); preprint OTIS-644, hep-ph/9712445.
5. H.Y. Cheng, preprint hep-ph/9712244; hep-ph/9807323; H.-Y. Cheng and B. Tseng, preprint IP-ASTP-01-98, hep-ph/9803457; hep-ph/9708211; hep-ph/9707316.
6. M. Bauer and B. Stech, Phys. Lett. **B152**, 380 (1985); M. Bauer, B. Stech and M. Wirbel, Z. Phys. **C34**, 103 (1987).
7. M. Neubert, B. Stech, CERN-TH-97-099, hep-ph/9705292, to appear in second edition of Heavy Flavours, ed. by A.J. Buras and M. Lindner, World Scientific, Singapore.
8. A. Ali, G. Kramer, C.-D. Lü, DESY-98-056, hep-ph/9805403, to appear in Phys. Rev. D .
9. R. Fleischer, T. Mannel, Phys. Rev. D57, 2752 (1998); preprint TTP-97-22, hep-ph/9706261; R. Fleischer, CERN-TH/98-254, hep-ph/9808238.